arXiv:hep-ph/0006199v1 16 Jun 2000

RIKEN-BNL preprint
June 2000

Prompt Photon Production in Polarized Hadron Collisions[†]

Werner Vogelsang[‡]

RIKEN-BNL Research Center, Brookhaven National Laboratory
Upton, NY 11973, U.S.A.

E-mail: wvogelsang@bnl.gov

Abstract

We consider spin asymmetries for prompt photon production in collisions of longitudinally polarized hadrons. This reaction will be a key tool at the BNL-RHIC $\vec{p}\vec{p}$ collider for determining the gluon spin density in a polarized proton. We study the effects of QCD corrections, such as all-order soft-gluon 'threshold' resummations.

[†] Talk presented at the "8th International Workshop on Deep-Inelastic Scattering" (DIS2000), 25-30 April 2000, Liverpool, UK

[‡] Thanks to RIKEN, Brookhaven National Laboratory and the U.S. Department of Energy (contract number DE-AC02-98CH10886) for providing the facilities essential for the completion of this work

PROMPT PHOTON PRODUCTION IN POLARIZED HADRON COLLISIONS

W. VOGELSANG
RIKEN-BNL Research Center, Brookhaven National Laboratory,
Upton, NY 11973, U.S.A.

We consider spin asymmetries for prompt photon production in collisions of longitudinally polarized hadrons. This reaction will be a key tool at the BNL-RHIC $\vec{p}\vec{p}$ collider for determining the gluon spin density in a polarized proton. We study the effects of QCD corrections, such as all-order soft-gluon 'threshold' resummations.

Prompt-photon production, $pp, p\bar{p}, pN \to \gamma X$, has been a classical tool for constraining the unpolarized gluon density. At leading order, a photon can be produced in the reactions $qg \to \gamma q$ and $q\bar{q} \to \gamma g$, giving rise to a distinct clean signal. Using *polarized* proton beams at RHIC offers a very promising method [1,2,3] to measure the *spin-dependent* gluon density in the nucleon, Δg, which is currently one of the most interesting quantities in nucleon structure.

A recent thorough theoretical next-to-leading order (NLO) QCD study [2] for prompt photon production at polarized RHIC also addressed the unwanted background from photons produced in jet fragmentation, when a parton, resulting from a QCD reaction, fragments into a photon plus a number of hadrons. In particular, the interplay between the fragmentation contribution and the chosen type of 'photon isolation cut', to be imposed in experiment [3] in order to reduce the background from π^0 decay photons, was studied. By comparing to lowest-order results, it was found in [2] that the QCD corrections to the polarized cross section are sizeable and reduce the dependence of the theory predictions on the factorization and renormalization scales; see Fig. 1.

In the unpolarized case, a pattern of disagreement between theoretical predictions and experimental data for prompt photon production has been observed in recent years [5,6,7]. The main problems reside in the fixed-target region, where NLO theory dramatically underpredicts some data sets [6,7]. At collider energies, as relevant to RHIC, there is less reason for concern, but also here the agreement is not satisfactory. Various improvements of the theoretical framework have been developed. One of them resorts to applying 'threshold' resummation to the prompt photon cross section [8], which organizes to all orders in α_s large logarithmic corrections to partonic hard scattering, associated with emission of soft gluons. It leads to a significant, albeit not sufficient, enhancement of the theory prediction in the fixed-target regime at large values of $p_T/\sqrt{s}$, accompanied by a dramatic reduction of scale dependence [9]. Thanks to the universal structure of soft-gluon emission, it is straightforward to apply



Figure 1: Left: polarized prompt photon cross section in $\vec{p}\vec{p}$ collisions at RHIC, at Born and NLO level, as function of photon transverse momentum. The upper half shows the scale dependence. The polarized parton densities were taken from [4]; for further details, see [2]. Right: spin asymmetry A_{LL} at LO, NLO, and including NLL threshold resummation.

threshold resummation to the polarized cross section. Fig. 1 shows the resulting effects on the *spin asymmetry* A_{LL}, for a 'toy' example that assumes a fictitious polarized set-up of the E706 experiment [6]. Details are as in [9]. Even though Fig. 1 does not directly refer to the case of RHIC, it is good news that resummation effects cancel to a large extent in A_{LL} for our present example.

Acknowledgments

I am grateful to S. Frixione and G. Sterman for fruitful collaborations.

References

1. E.L. Berger and J. Qiu, *Phys. Rev.* D **40**, 778 (1989).
2. S. Frixione and W. Vogelsang, *Nucl. Phys.* B **568**, 60 (2000).
3. L.C. Bland, `hep-ex/9907058`;
   Y. Goto, *Nucl. Phys. Proc. Suppl.* **79**, 588 (1999).
4. M. Glück *et al.*, *Phys. Rev.* D **53**, 4775 (96).
5. F. Abe et al., CDF Collab., *Phys. Rev. Lett.* **73**, 2662 (1994).
6. L. Apanasevich et al., E706 Collab., *Phys. Rev. Lett.* **81**, 2642 (1998).
7. G. Ballocchi et al., UA6 Collab., *Phys. Lett.* B **436**, 222 (1998).
8. E. Laenen, G. Oderda, G. Sterman, *Phys. Lett.* B **438**, 173 (1998);
   S. Catani, M.L. Mangano, P. Nason, *JHEP* **9807**, 024 (1998);
   N. Kidonakis and J.F. Owens, *Phys. Rev.* D **61**, 094004 (2000).
9. S. Catani, M.L. Mangano, P. Nason, C. Oleari, W. Vogelsang,
   JHEP **9903**, 025 (1999).